

BREED MIX



GENETIC STATS


Wolfiness: 1.6 % **HIGH**
Predicted adult weight: **25 lbs**
Life stage: **Young adult**
Based on your dog's date of birth provided.

TEST DETAILS


Kit number: EM-72185494
Swab number: 31241260135269

BREED MIX BY CHROMOSOME

Our advanced test identifies from where Zari inherited every part of the chromosome pairs in her genome.



FAMILY TREE





(<https://app.embarkvet.com/pet/9cf48ca4-d791-4ad3-abf9-29056adb832d/breed-reveal>)



SIBERIAN HUSKY



The Siberian Husky originated from the extreme north east of Siberia. They were initially domesticated by the Chukchi -an ancient population that thrived by herding reindeer and moving with each season to new grazing regions. They came to America in 1909 and found their place in the Alaskan wilderness. They love to be out in cold weather and are known to be the ideal sled dog. They have strong insulated paws that are perfect for traction in the snow. The Siberian Husky also has two layers in their coat that protects them from Arctic winters.

Fun Fact

In 1925 a team of Siberian Huskies saved Nome, Alaska by carrying the serum to cure diphtheria a considerable distance by sled. The run was done in the middle of a blizzard and in conditions below -23 degrees Fahrenheit. The run is remembered by the annual Iditarod Trail Sled Race, and Balto, the famous sled dog who led his team through the final leg.

POMERANIAN



Fun Fact

Pomeranians boast one of the widest variety of color options in one breed. The American Kennel Club lists 23 accepted colors

Cute, feisty and furry, Poms are intelligent and loyal to their families. Don't let their cuteness fool you, however. These independent, bold dogs have minds of their own. They are alert and curious about the world around them. Unfortunately, in their minds, they are much larger than they really are, which can sometimes lead them to harass and even attack much larger dogs. Luckily, if they are properly socialized with other dogs and animals, they generally get along quite well with them. Poms take their name from the province of Pomerania, in Germany. They became especially popular when Queen Victoria allowed some of her Pomeranians to be shown in a conformation show, the first Pomeranians ever to be shown. Pomeranians make excellent pets for older people and those who are busy, because they aren't an overly dependent breed. They are also good for apartment dwellers or homes that don't have a backyard. Because of their small size, they aren't recommended for families with small children who might injure them accidentally. While Poms are good with children, they are not a good choice for very young or highly active children because of their small size. Never let your small children and your Pom play without supervision. Because they are so small, Poms can be perceived as prey by owls, eagles, hawks, coyotes, and other wild animals. Never leave them outside unattended, and be watchful if there are predatory birds in your location. If this is the case, stay close to your Pom to discourage birds from trying to carry them off!

AMERICAN ESKIMO DOG



The American Eskimo Dog, or Eskie, is a fluffy little fellow with a misleading name. The breed actually originated in Germany and is likely a close descendent of the German Spitz. These guys gained popularity in America during the 1800's while serving as circus dogs. Eskies are highly intelligent, but they take much longer to mature and leave that puppy behavior behind. The breed comes in three different sizes: toy, miniature, and standard. Their fluffy white double coat requires regular brushing and grooming. With good training, and adequate exercise American Eskimo Dogs can make the best of companions!

Fun Fact

The American Eskimo Dog used to be called the American Spitz, but it is believed that a kennel renamed the breed using their own name, American Eskimo Kennel.

MATERNAL LINE



Through Zari's mitochondrial DNA we can trace her mother's ancestry back to where dogs and people first became friends. This map helps you visualize the routes that her ancestors took to your home. Their story is described below the map.

HAPLOGROUP: C1

Congratulations, C1 is a very exotic female lineage! It is more closely associated with maternal lineages found in wolves, foxes and jackals than with other dog lineages. So it seems dogs in this group have a common male dog ancestor who, many thousands of years ago, mated with a female wolf! This is not a common lineage in any breed, though a good number of German Shepherds and Doberman Pinchers are C1. It is also found in breeds as diverse as Peruvian Inca Orchids and Pekingese; it is rarely found amongst Labrador Retrievers, Border Collies, Siberian Huskies, or Cocker Spaniels. Despite its fascinating origins, it is widely distributed around the globe, and even shows up frequently among Peruvian village dogs. It almost certainly survived at low frequency in Europe for millennia and then was dispersed outside of Europe by colonialism, though not as successfully as some other lineages.

HAPLOTYPE: C39

Part of the C1 haplogroup, this haplotype occurs most frequently in Pomeranians and Xoloitzcuintli.

TRAITS: BASE COAT COLOR

TRAIT

RESULT

Dark or Light Fur | *E (Extension) Locus* | Gene: *Melanocortin Receptor 1 (MC1R)* | Genetic Result: **E^ee**

This gene helps determine whether a dog can produce dark (black or brown) hairs or lighter yellow or red hairs. Any result except for **ee** means that the dog can produce dark hairs. An **ee** result means that the dog does not produce dark hairs and will have lighter yellow or red hairs all over its entire body.

The overall MC1R genetic result is influenced by more subloci than those presented in this section. Additional MC1R subloci results can be found under the **Coat Color Modifiers > Facial Fur Pattern** section below.

Did You Know? If a dog has an **ee** result, then the fur's actual shade can range from a deep copper to white - the exact color cannot be predicted solely from this result and will depend on other genetic factors, including the red pigment intensity test.

Can have dark fur

Dark brown pigment | *Cocoa* | Gene: *HPS3* | Genetic Result: **NN**

Dogs with the **coco** genotype will produce dark brown pigment instead of black in both their hair and skin. Dogs with the **Nco** genotype will produce black pigment, but can pass the **co** variant on to their puppies. Dogs that have the **coco** genotype as well as the **bb** genotype at the B locus are generally a lighter brown than dogs that have the **Bb** or **BB** genotypes at the B locus.

Did You Know? The **co** variant and the dark brown "cocoa" coat color have only been documented in French Bulldogs. Dogs with the cocoa coat color are sometimes born with light brown coats that darken as they reach maturity.

No impact on fur and skin color

Red Pigment Intensity | *I (Intensity) Loci* | Genetic Result: **Dilute Red Pigmentation**

Intensity refers to the concentration of red pigment in the coat. Dogs with more densely concentrated (intense) pigment will be a deeper red, while dogs with less concentrated (dilute) pigment will be tan, yellow, cream, or white. Five locations in the dog genome explain approximately 70% of red pigmentation intensity variation across all dogs. Because the locations we test may not directly cause differences in red pigmentation intensity, we consider this to be a linkage test.

Did You Know? One of the genes that influences pigment intensity in dogs, TYR, is also responsible for intensity variation in domestic mice, cats, cattle, rabbits, and llamas. In dogs and humans, more genes are involved.

Any light fur likely white or cream

TRAITS: BASE COAT COLOR (CONTINUED)

TRAIT

RESULT

Brown or Black Pigment | *B (Brown) Locus* | *Gene: Tyrosinase Related Protein 1 (TYRP1)* | Genetic Result: **BB**

This gene helps determine whether a dog produces brown or black pigments. Dogs with a **bb** result produce brown pigment instead of black in both their hair and skin, while dogs with a **Bb** or **BB** result produce black pigment. Dogs that have **ee** at the E (Extension) Locus and **bb** at this B (Brown) Locus are likely to have red or cream coats and brown noses, eye rims, and footpads, which is sometimes referred to as "Dudley Nose" in Labrador Retrievers.

Black or gray fur and skin

Did You Know? "Liver" or "chocolate" is the preferred color term for brown in most breeds; in the Doberman Pinscher it is referred to as "red".

Color Dilution | *D (Dilute) Locus* | *Gene: Melanophilin (MLPH)* | Genetic Result: **DD**

This gene helps determine whether a dog has lighter "diluted" pigment. A dog with a **Dd** or **DD** result will not be dilute. A dog with a **dd** result will have all their black or brown pigment lightened ("diluted") to gray or light brown, and may lighten red pigment to cream. This affects their fur, skin, and sometimes eye color. The D locus result that we report is determined by three different genetic variants that can work together to cause diluted pigmentation. These are the common **d** allele, also known as "**d1**", and the less common alleles known as "**d2**" and "**d3**". Dogs with two **d** alleles, regardless of which variant, are typically dilute.

Dark (non-dilute) fur and skin

Did You Know? There are many breed-specific names for these dilute colors, such as "blue", "charcoal", "fawn", "silver", and "Isabella". Dilute dogs, especially in certain breeds, have a higher incidence of Color Dilution Alopecia which causes hair loss in some patches.

TRAITS: COAT COLOR MODIFIERS

TRAIT

RESULT

Hidden Patterning | *K (Dominant Black) Locus* | Gene: *Canine Beta-Defensin 103 (CBD103)* | Genetic Result: **k^Yk^Y**

This gene helps determine whether the dog has a black coat. Dogs with a **k^Yk^Y** result will show a coat color pattern based on the result they have at the A (Agouti) Locus. A **K^BK^B** or **K^Bk^Y** result means the dog is dominant black, which overrides the fur pattern that would otherwise be determined by the A (Agouti) Locus. These dogs will usually have solid black or brown coats, or if they have **ee** at the E (Extension) Locus then red/cream coats, regardless of their result at the A (Agouti) Locus. Dogs who test as **K^Bk^Y** may be brindle rather than black or brown.

More likely to have patterned fur

Did You Know? Even if a dog is "dominant black" several other genes could still impact the dog's fur and cause other patterns, such as white spotting.

Body Pattern | *A (Agouti) Locus* | Gene: *Agouti Signalling Protein (ASIP)* | Genetic Result: **a^wa^w**

This gene is responsible for causing different coat patterns. It only affects the fur of dogs that do not have **ee** at the E (Extension) Locus and do have **k^Yk^Y** at the K (Dominant Black) Locus. It controls switching between black and red pigment in hair cells, which means that it can cause a dog to have hairs that have sections of black and sections of red/cream, or hairs with different colors on different parts of the dog's body. Sable or Fawn dogs have a mostly or entirely red coat with some interspersed black hairs. Agouti or Wolf Sable dogs have red hairs with black tips, mostly on their head and back. Black and tan dogs are mostly black or brown with lighter patches on their cheeks, eyebrows, chest, and legs. Recessive black dogs have solid-colored black or brown coats.

Agouti (Wolf Sable) coat color pattern

Did You Know? The ASIP gene causes interesting coat patterns in many other species of animals as well as dogs.

TRAITS: COAT COLOR MODIFIERS (CONTINUED)

TRAIT

RESULT

Facial Fur Pattern | *E (Extension) Locus* | Gene: *Melanocortin Receptor 1 (MC1R)* | Genetic Result: **E^me**

This gene determines whether a dog can have dark hair and can give it a black "mask" or "widow's peak," unless the dog has overriding coat color genetic factors. Dogs with one or two copies of **E^m** in their result may have a mask, which is dark facial fur as seen in the German Shepherd Dog and Pug. Dogs with no **E^m** in their result but one or two copies of the **E^g**, **E^a**, or **E^h** variants can instead have a "widow's peak", which is dark forehead fur.

Did You Know?

The "widow's peak" is seen in the Afghan Hound and Borzoi, and is called either "grizzle" or "domino."

In the absence of **E^m**, dogs with the **E^g** variant can have a "widow's peak" phenotype. In the absence of both **E^m** and **E** variants, dogs with the **E^a** or **E^h** variants can express the "widow's peak" phenotype. Additionally, a dog with any combination of two of the **E^g**, **E^a**, or **E^h** variants (example: **E^gE^a**) is also expected to express the grizzle phenotype.

Can have
grizzle/domino
"Widow's Peak" (dark
forehead fur)

Saddle Tan | Gene: *RALY* | Genetic Result: **NN**

The "Saddle Tan" pattern causes the black hairs to recede into a "saddle" shape on the back, leaving a tan face, legs, and belly, as a dog ages. The Saddle Tan pattern is characteristic of breeds like the Corgi, Beagle, and German Shepherd. Dogs that have the **ll** genotype at this locus are more likely to be mostly black with tan points on the eyebrows, muzzle, and legs as commonly seen in the Doberman Pinscher and the Rottweiler. This gene modifies the A Locus **a^t** allele, so dogs that do not express **a^t** are not influenced by this gene.

Did You Know? The Saddle Tan pattern is characteristic of breeds like the Corgi, Beagle, and German Shepherd.

No impact on coat
pattern

TRAITS: COAT COLOR MODIFIERS (CONTINUED)

TRAIT

RESULT

White Spotting | *S (White Spotting) Locus* | *Gene: MITF* | Genetic Result: **spsp**

This gene is responsible for most of the white spotting observed in dogs. Dogs with a result of **spsp** will have a nearly white coat or large patches of white in their coat. Dogs with a result of **Ssp** will have more limited white spotting that is breed-dependent. A result of **SS** means that a dog likely has no white or minimal white in their coat. The S Locus does not explain all white spotting patterns in dogs and other causes are currently being researched. Some dogs may have small amounts of white on the paws, chest, face, or tail regardless of their result at this gene.

Likely to have large white areas in coat

Did You Know? Any dog can have white spotting regardless of coat color. The colored sections of the coat will reflect the dog's other genetic coat color results.

Roan | *R (Roan) Locus* | *Gene: USH2A* | Genetic Result: **rr**

This gene, along with the S Locus, regulates whether a dog will have roaning. Dogs with at least one copy of **R** will likely have roaning on otherwise uniformly unpigmented white areas created by the S Locus. Roan may not be visible if white spotting is limited to small areas, such as the paws, chest, face, or tail. The extent of roaning varies from uniform roaning to non-uniform roaning, and patchy, non-uniform roaning may look similar to ticking. Roan does not appear in white areas created by other genes, such as a combination of the E Locus and I Locus (for example, Samoyeds). The roan pattern can appear with or without ticking.

Likely no impact on coat pattern

Did You Know? Roan, tick, and Dalmatians' spots become visible a few weeks after birth. The R Locus is probably involved in the development of Dalmatians' spots.

Merle | *M (Merle) Locus* | *Gene: PMEL* | Genetic Result: **mm**

This gene is responsible for mottled or patchy coat color in some dogs. Dogs with an **M*m** result are likely to appear merle or could be "non-expressing" merle, meaning that the merle pattern is very subtle or not at all evident in their coat. Dogs with an **M*M*** result are likely to have merle or double merle coat patterning. Dogs with an **mm** result are unlikely to have a merle coat pattern.

Unlikely to have merle pattern

Did You Know? Merle coat patterning is common to several dog breeds including the Australian Shepherd, Catahoula Leopard Dog, and Shetland Sheepdog.

TRAITS: COAT COLOR MODIFIERS (CONTINUED)

TRAIT

RESULT

Harlequin | Gene: *PSMB* | Genetic Result: **hh**

This gene, along with the M Locus, determines whether a dog will have harlequin patterning. This pattern is recognized in Great Danes and causes dogs to have a white coat with patches of darker pigment. A dog with an **Hh** result will be harlequin if they are also **M*m** or **M*M*** at the M Locus and are not **ee** at the E locus. Dogs with a result of **hh** will not be harlequin.

No impact on coat pattern

Did You Know? While many harlequin dogs are white with black patches, some dogs have grey, sable, or brindle patches of color, depending on their genotypes at other coat color genes.

Panda White Spotting | Gene: *KIT* | Genetic Result: **NN**

Panda White Spotting originated in a line of German Shepherd Dogs and causes a mostly symmetrical white spotting of the head and/or body. This is a dominant variant of the KIT gene, which has a role in pigmentation.

Dogs with one copy of the **I** allele will exhibit this white spotting. Dogs with two copies of the **I** allele have never been observed, as two copies of the variant is suspected to be lethal to the developing embryo.

Dogs with the **NN** result will not exhibit white spotting due to this variant.

Not expected to display Panda pattern

Did You Know? A de novo mutation (a genetic mutation not inherited from the parents) occurred in a female German Shepherd Dog named Lewcinka's Franka von Phenom. She was born in 2000, and all Panda Shepherds can trace their bloodline back to her.

TRAITS: OTHER COAT TRAITS

TRAIT

RESULT

Furnishings | Gene: *RSPO2* | Genetic Result: **II**

This gene is responsible for “furnishings”, which is another name for the mustache, beard, and eyebrows that are characteristic of breeds like the Schnauzer, Scottish Terrier, and Wire Haired Dachshund. A dog with an **FF** or **FI** result is likely to have furnishings. A dog with an **II** result will not have furnishings. We measure this result using a linkage test.

Likely unfurnished (no mustache, beard, and/or eyebrows)

Did You Know? In breeds that are expected to have furnishings, dogs without furnishings are the exception - this is sometimes called an “improper coat”.

Coat Length | Gene: *FGF5* | Genetic Result: **LhLh**

This gene affects hair length in many species, including cats, dogs, mice, and humans. In dogs, an **Lh** allele confers a long, silky hair coat across many breeds, including Yorkshire Terriers, Cocker Spaniels, and Golden Retrievers. An **ShSh** or **ShLh** result is likely to mean a shorter coat, like in the Boxer or the American Staffordshire Terrier. The coat length determined by *FGF5*, as reported by us, is influenced by four genetic variants that work together to promote long hair.

The most common of these is the **Lh1** variant (G/T, CanFam3.1, chr32, g.4509367) and the less common ones are **Lh2** (C/T, CanFam3.1, chr32, g.4528639), **Lh3** (16bp deletion, CanFam3.1, chr32, g.4528616), and **Lh4** (GG insertion, CanFam3.1, chr32, g.4528621). The *FGF5_Lh1* variant is found across many dog breeds. The less common variants, *FGF5_Lh2* have been found in the Akita, Samoyed, and Siberian Husky, *FGF5_Lh3* have been found in the Eurasier, and *FGF5_Lh4* have been found in the Afghan Hound, Eurasier, and French Bulldog.

Likely long coat

The **Lh** alleles have a recessive mode of inheritance, meaning that two copies of the **Lh** alleles are required to have long hair. The presence of two *Lh* alleles at any of these *FGF5* loci is expected to result in long hair. One copy each of **Lh1** and **Lh2** have been found in Samoyeds, one copy each of **Lh1** and **Lh3** have been found in Eurasiers and one copy each of **Lh1** and **Lh4** have been found in Afghan Hounds and Eurasiers.

Did You Know? In certain breeds, such as Pembroke Welsh Corgi and French Bulldog, the long coat is described as “fluffy.”

TRAITS: OTHER COAT TRAITS (CONTINUED)

| TRAIT | RESULT |
|--|--|
| <p>Shedding <i>Gene: MC5R</i> Genetic Result: CC</p> <p>This gene affects how much a dog sheds. Dogs with furnishings or wire-haired coats tend to be low shedders regardless of their result for this gene. In other dogs, a CC or CT result indicates heavy or seasonal shedding, like many Labradors and German Shepherd Dogs. Dogs with a TT result tend to be lighter shedders, like Boxers, Shih Tzus and Chihuahuas.</p> | <p>Likely heavy/seasonal shedding</p> |
| <p>Coat Texture <i>Gene: KRT71</i> Genetic Result: CC</p> <p>For dogs with long fur, dogs with a TT or CT result will likely have a wavy or curly coat like the coat of Poodles and Bichon Frises. Dogs with a CC result will likely have a straight coat—unless the dog has a "Likely Furnished" result for the Furnishings trait, since this can also make the coat more curly.</p> <p>Did You Know? Dogs with short coats may have straight coats, whatever result they have for this gene.</p> | <p>Likely straight coat</p> |
| <p>Hairlessness (Xolo type) <i>Gene: FOXI3</i> Genetic Result: NN</p> <p>This gene can cause hairlessness over most of the body as well as changes in tooth shape and number. This particular gene occurs in Peruvian Inca Orchid, Xoloitzcuintli (Mexican Hairless), and Chinese Crested; other hairless breeds are due to different genes. Dogs with the NDup result are likely to be hairless while dogs with the NN result are likely to have a normal coat. We measure this result using a linkage test.</p> <p>Did You Know? The DupDup result has never been observed, suggesting that dogs with that genotype cannot survive to birth.</p> | <p>Very unlikely to be hairless</p> |
| <p>Hairlessness (Terrier type) <i>Gene: SGK3</i> Genetic Result: NN</p> <p>This gene is responsible for Hairlessness in the American Hairless Terrier. Dogs with the DD result are likely to be hairless. Dogs with the ND genotype will have a normal coat, but can pass the D variant on to their offspring.</p> | <p>Very unlikely to be hairless</p> |

Oculocutaneous Albinism Type 2 | Gene: *SLC45A2* | Genetic Result: **NN**

This gene causes oculocutaneous albinism (OCA), also known as Doberman Z Factor Albinism. Dogs with a **DD** result will have OCA. Effects include severely reduced or absent pigment in the eyes, skin, and hair, and sometimes vision problems due to lack of eye pigment (which helps direct and absorb ambient light) and are prone to sunburn. Dogs with a **ND** result will not be affected, but can pass the mutation on to their offspring. We measure this result using a linkage test.

Likely not albino

Did You Know? This particular mutation can be traced back to a single white Doberman Pinscher born in 1976, and it has only been observed in dogs descended from this individual.



TRAITS: OTHER BODY FEATURES

TRAIT

RESULT

Muzzle Length | Gene: *BMP3* | Genetic Result: **CC**

This gene affects muzzle length. A dog with a **AC** or **CC** result is likely to have a medium-length muzzle like a Staffordshire Terrier or Labrador, or a long muzzle like a Whippet or Collie. A dog with a **AA** result is likely to have a short muzzle, like an English Bulldog, Pug, or Pekingese.

Did You Know? At least five different genes affect snout length in dogs, with *BMP3* being the only one with a known causal mutation. For example, the muzzle length of some breeds, including the long-snouted Scottish Terrier or the short-snouted Japanese Chin, appear to be caused by other genes. This means your dog may have a long or short snout due to other genetic factors. Embark is working to figure out what these might be.

Likely medium or long muzzle

Tail Length | Gene: *T* | Genetic Result: **CC**

This is one of the genes that can cause a short bobtail. Most dogs have a **CC** result and a long tail. Dogs with a **CG** result are likely to have a bobtail, which is an unusually short or absent tail. This can be seen in many "natural bobtail" breeds including the Pembroke Welsh Corgi, the Australian Shepherd, and the Brittany Spaniel. Dogs with **GG** genotypes have not been observed, suggesting that dogs with such a result do not survive to birth.

Did You Know? While certain lineages of Boston Terrier, English Bulldog, Rottweiler, Miniature Schnauzer, Cavalier King Charles Spaniel, and Parson Russell Terrier, and Dobermans are born with a natural bobtail, it is not always caused by this gene. This suggests that other unknown genetic effects can also lead to a natural bobtail.

Likely normal-length tail

Hind Dew Claws | Gene: *LMBR1* | Genetic Result: **CT**

This is one of the genes that can cause hind dew claws, which are extra, nonfunctional digits located midway between a dog's paw and hock. Dogs with a **CT** or **TT** result have about a 50% chance of having hind dewclaws. Hind dew claws can also be caused by other, still unknown, genes. Embark is working to figure those out.

Did You Know? Hind dew claws are commonly found in certain breeds such as the Saint Bernard.

Likely to have hind dew claws

TRAITS: OTHER BODY FEATURES (CONTINUED)

TRAIT

RESULT

Back Muscling & Bulk (Large Breed) | Gene: *ACSL4* | Genetic Result: **CC**

This gene can cause heavy muscling along the back and trunk in characteristically "bulky" large-breed dogs including the Saint Bernard, Bernese Mountain Dog, Greater Swiss Mountain Dog, and Rottweiler. A dog with the **TT** result is likely to have heavy muscling. Leaner-shaped large breed dogs like the Great Dane, Irish Wolfhound, and Scottish Deerhound generally have a **CC** result. The **TC** result also indicates likely normal muscling.

Likely normal muscling

Did You Know? This gene does not seem to affect muscling in small or even mid-sized dog breeds with lots of back muscling, including the American Staffordshire Terrier, Boston Terrier, and the English Bulldog.

Eye Color | Gene: *ALX4* | Genetic Result: **NN**

This gene is associated with blue eyes in Arctic breeds like Siberian Husky as well as tri-colored (non-merle) Australian Shepherds. Dogs with a **DupDup** or **NDup** result are more likely to have blue eyes, although some dogs may have only one blue eye or may not have blue eyes at all; nevertheless, they can still pass blue eyes to their offspring. Dogs with a **NN** result may have blue eyes due to other factors, such as merle or white spotting. We measure this result using a linkage test.

Less likely to have blue eyes

Did You Know? Embark researchers discovered this gene by studying data from dogs like yours. Who knows what we will be able to discover next? Answer the questions on our research surveys to contribute to future discoveries!

Chondrodysplasia (Leg Length) | Gene: *Chr. 18 FGF4 Retrogene* | Genetic Result: **NN**

This variant is associated with a type of disproportionate dwarfism known as chondrodysplasia (CDPA). CDPA is a breed-defining characteristic of many breeds exhibiting a "short-legged, long-bodied" appearance, such as Corgis, Dachshunds, Basset Hounds, and others. Dogs with the **II** result display the largest reduction in leg length. Dogs with the **NI** genotype will have an intermediate leg length, while dogs with the **NN** result will not exhibit leg shortening due to this variant.

Likely to have normal leg length

Did You Know? A similar genetic variant called the chondrodystrophy (CDDY) variant also plays an important role in shortening the leg length of many breeds. Dog breeds with the shortest legs, like the Corgi, Dachshund, and Basset Hound generally have one or two copies of the CDDY and CDPA variants. CDDY (but not CDPA) is also associated with an increased risk of Type I Intervertebral Disc Disease (IVDD). You can see the CDDY result in the health test results under "Intervertebral Disc Disease Type I".

TRAITS: BODY SIZE

| TRAIT | RESULT |
|--|---------------------|
| Body Size 1 <i>Gene: IGF1</i> Genetic Result: NI This is one of several genes that influence the size of a dog. A result of II for this gene is associated with smaller body size. A result of NN is associated with larger body size. | Intermediate |
| Body Size 2 <i>Gene: IGFR1</i> Genetic Result: GG This is one of several genes that influence the size of a dog. A result of AA for this gene is associated with smaller body size. A result of GG is associated with larger body size. | Larger |
| Body Size 3 <i>Gene: STC2</i> Genetic Result: TA This is one of several genes that influence the size of a dog. A result of AA for this gene is associated with smaller body size. A result of TT is associated with larger body size. | Intermediate |
| Body Size 4 <i>Gene: GHR - E191K</i> Genetic Result: AA This is one of several genes that influence the size of a dog. A result of AA for this gene is associated with smaller body size. A result of GG is associated with larger body size. | Smaller |
| Body Size 5 <i>Gene: GHR - P177L</i> Genetic Result: CC This is one of several genes that influence the size of a dog. A result of TT for this gene is associated with smaller body size. A result of CC is associated with larger body size. | Larger |

TRAITS: PERFORMANCE

TRAIT

RESULT

Altitude Adaptation | Gene: *EPAS1* | Genetic Result: **GG**

This gene causes dogs to be especially tolerant of low oxygen environments, such as those found at high elevations. Dogs with a **AA** or **GA** result will be less susceptible to "altitude sickness."

Normal altitude tolerance

Did You Know? This gene was originally identified in breeds from high altitude areas such as the Tibetan Mastiff.

Appetite | Gene: *POMC* | Genetic Result: **NN**

This gene influences eating behavior. An **ND** or **DD** result would predict higher food motivation compared to **NN** result, increasing the likelihood to eat excessively, have higher body fat percentage, and be more prone to obesity. Read more about the genetics of POMC, and learn how you can contribute to research, in our blog post (<https://embarkvet.com/resources/blog/pomc-dogs/>). We measure this result using a linkage test.

Normal food motivation

Did You Know? POMC is actually short for "proopiomelanocortin," and is a large protein that is broken up into several smaller proteins that have biological activity. The smaller proteins generated from POMC control, among other things, distribution of pigment to the hair and skin cells, appetite, and energy expenditure.

HEALTH REPORT

How to interpret Zari's genetic health results:

If Zari inherited any of the variants that we tested, they will be listed at the top of the Health Report section, along with a description of how to interpret this result. We also include all of the variants that we tested Zari for that we did not detect the risk variant for.

A genetic test is not a diagnosis

This genetic test does not diagnose a disease. Please talk to your vet about your dog's genetic results, or if you think that your pet may have a health condition or disease.

Summary

Of the 274 genetic health risks we analyzed, we found 2 results that you should learn about.

Notable results (2)

ALT Activity

Degenerative Myelopathy, DM














Clear results

Breed-relevant (12)

Other (259)



















BREED-RELEVANT RESULTS

Research studies indicate that these results are more relevant to dogs like Zari, and may influence her chances of developing certain health conditions.



















| | |
|---|---------|
|  Degenerative Myelopathy, DM (SOD1A) | Notable |
|  Day Blindness (CNGB3 Deletion, Alaskan Malamute Variant) | Clear |
|  GM1 Gangliosidosis (GLB1 Exon 15, Alaskan Husky Variant) | Clear |
|  Hereditary Vitamin D-Resistant Rickets (VDR) | Clear |
|  Methemoglobinemia (CYB5R3) | Clear |
|  Oculocutaneous Albinism, OCA (SLC45A2, Small Breed Variant) | Clear |
|  Primary Lens Luxation (ADAMTS17) | Clear |
|  Progressive Retinal Atrophy 5, PRA5 (NECAP1 Exon 6, Giant Schnauzer Variant) | Clear |
|  Progressive Retinal Atrophy, prcd (PRCD Exon 1) | Clear |
|  Progressive Retinal Atrophy, rcd3 (PDE6A) | Clear |
|  Thrombopathia (RASGRP1 Exon 5, American Eskimo Dog Variant) | Clear |
|  Urate Kidney & Bladder Stones (SLC2A9) | Clear |
|  X-Linked Progressive Retinal Atrophy 1, XL-PRA1 (RPGR) | Clear |

OTHER RESULTS



















Research has not yet linked these conditions to dogs with similar breeds to Zari. Review any increased risk or notable results to understand her potential risk and recommendations.

| | |
|--|---------|
|  ALT Activity (GPT) | Notable |
|  2-DHA Kidney & Bladder Stones (APRT) | Clear |
|  Acral Mutilation Syndrome (GDNF-AS, Spaniel and Pointer Variant) | Clear |
|  Alaskan Husky Encephalopathy (SLC19A3) | Clear |
|  Alaskan Malamute Polyneuropathy, AMPN (NDRG1 SNP) | Clear |
|  Alexander Disease (GFAP) | Clear |
|  Anhidrotic Ectodermal Dysplasia (EDA Intron 8) | Clear |
|  Autosomal Dominant Progressive Retinal Atrophy (RHO) | Clear |
|  Bald Thigh Syndrome (IGFBP5) | Clear |
|  Bernard-Soulier Syndrome, BSS (GP9, Cocker Spaniel Variant) | Clear |
|  Bully Whippet Syndrome (MSTN) | Clear |
|  Canine Elliptocytosis (SPTB Exon 30) | Clear |
|  Canine Fucosidosis (FUCA1) | Clear |
|  Canine Leukocyte Adhesion Deficiency Type I, CLAD I (ITGB2, Setter Variant) | Clear |
|  Canine Leukocyte Adhesion Deficiency Type III, CLAD III (FERMT3, German Shepherd Variant) | Clear |
|  Canine Multifocal Retinopathy, cmr1 (BEST1 Exon 2) | Clear |
|  Canine Multifocal Retinopathy, cmr2 (BEST1 Exon 5, Coton de Tulear Variant) | Clear |
|  Canine Multifocal Retinopathy, cmr3 (BEST1 Exon 10 Deletion, Finnish and Swedish Lapphund, Lapponian Herder Variant) | Clear |



















OTHER RESULTS

| | |
|--|-------|
|  Canine Multiple System Degeneration (SERAC1 Exon 4, Chinese Crested Variant) | Clear |
|  Canine Multiple System Degeneration (SERAC1 Exon 15, Kerry Blue Terrier Variant) | Clear |
|  Cardiomyopathy and Juvenile Mortality (YARS2) | Clear |
|  Centronuclear Myopathy, CNM (PTPLA) | Clear |
|  Cerebellar Hypoplasia (VLDLR, Eurasier Variant) | Clear |
|  Chondrodysplasia (ITGA10, Norwegian Elkhound and Karelian Bear Dog Variant) | Clear |
|  Cleft Lip and/or Cleft Palate (ADAMTS20, Nova Scotia Duck Tolling Retriever Variant) | Clear |
|  Cleft Palate, CP1 (DLX6 intron 2, Nova Scotia Duck Tolling Retriever Variant) | Clear |
|  Cobalamin Malabsorption (CUBN Exon 8, Beagle Variant) | Clear |
|  Cobalamin Malabsorption (CUBN Exon 53, Border Collie Variant) | Clear |
|  Collie Eye Anomaly (NHEJ1) | Clear |
|  Complement 3 Deficiency, C3 Deficiency (C3) | Clear |
|  Congenital Cornification Disorder (NSDHL, Chihuahua Variant) | Clear |
|  Congenital Dyserythropoietic Anemia and Polymyopathy (EHPB1L1, Labrador Retriever Variant) | Clear |
|  Congenital Hypothyroidism (TPO, Rat, Toy, Hairless Terrier Variant) | Clear |
|  Congenital Hypothyroidism (TPO, Tenterfield Terrier Variant) | Clear |
|  Congenital Hypothyroidism with Goiter (TPO Intron 13, French Bulldog Variant) | Clear |
|  Congenital Hypothyroidism with Goiter (SLC5A5, Shih Tzu Variant) | Clear |



















OTHER RESULTS

| | |
|--|-------|
|  Congenital Macrothrombocytopenia (TUBB1 Exon 1, Cairn and Norfolk Terrier Variant) | Clear |
|  Congenital Muscular Dystrophy (LAMA2, Italian Greyhound) | Clear |
|  Congenital Myasthenic Syndrome, CMS (COLQ, Labrador Retriever Variant) | Clear |
|  Congenital Myasthenic Syndrome, CMS (COLQ, Golden Retriever Variant) | Clear |
|  Congenital Myasthenic Syndrome, CMS (CHAT, Old Danish Pointing Dog Variant) | Clear |
|  Congenital Myasthenic Syndrome, CMS (CHRNE, Jack Russell Terrier Variant) | Clear |
|  Congenital Stationary Night Blindness (LRIT3, Beagle Variant) | Clear |
|  Congenital Stationary Night Blindness (RPE65, Briard Variant) | Clear |
|  Copper Toxicosis (Accumulating) (ATP7B) | Clear |
|  Copper Toxicosis (Attenuating) (ATP7A, Labrador Retriever) | Clear |
|  Copper Toxicosis (Attenuating) (RETN, Labrador Retriever) | Clear |
|  Craniomandibular Osteopathy, CMO (SLC37A2) | Clear |
|  Craniomandibular Osteopathy, CMO (SLC37A2 Intron 16, Basset Hound Variant) | Clear |
|  Cystinuria Type I-A (SLC3A1, Newfoundland Variant) | Clear |
|  Cystinuria Type II-A (SLC3A1, Australian Cattle Dog Variant) | Clear |
|  Cystinuria Type II-B (SLC7A9, Miniature Pinscher Variant) | Clear |
|  Darier Disease (ATP2A2, Irish Terrier Variant) | Clear |
|  Day Blindness (CNGA3 Exon 7, German Shepherd Variant) | Clear |

OTHER RESULTS

| | |
|--|-------|
|  Day Blindness (CNGA3 Exon 7, Labrador Retriever Variant) | Clear |
|  Day Blindness (CNGB3 Exon 6, German Shorthaired Pointer Variant) | Clear |
|  Deafness and Vestibular Syndrome of Dobermans, DVDob, DINGS (MYO7A) | Clear |
|  Demyelinating Polyneuropathy (SBF2/MTRM13) | Clear |
|  Dental-Skeletal-Retinal Anomaly (MIA3, Cane Corso Variant) | Clear |
|  Diffuse Cystic Renal Dysplasia and Hepatic Fibrosis (INPP5E Intron 9, Norwich Terrier Variant) | Clear |
|  Dilated Cardiomyopathy, DCM (RBM20, Schnauzer Variant) | Clear |
|  Dilated Cardiomyopathy, DCM1 (PDK4, Doberman Pinscher Variant 1) | Clear |
|  Dilated Cardiomyopathy, DCM2 (TTN, Doberman Pinscher Variant 2) | Clear |
|  Disproportionate Dwarfism (PRKG2, Dogo Argentino Variant) | Clear |
|  Dry Eye Curly Coat Syndrome (FAM83H Exon 5) | Clear |
|  Dystrophic Epidermolysis Bullosa (COL7A1, Central Asian Shepherd Dog Variant) | Clear |
|  Dystrophic Epidermolysis Bullosa (COL7A1, Golden Retriever Variant) | Clear |
|  Early Bilateral Deafness (LOXHD1 Exon 38, Rottweiler Variant) | Clear |
|  Early Onset Adult Deafness, EOAD (EPS8L2 Deletion, Rhodesian Ridgeback Variant) | Clear |
|  Early Onset Cerebellar Ataxia (SEL1L, Finnish Hound Variant) | Clear |
|  Ehlers Danlos (ADAMTS2, Doberman Pinscher Variant) | Clear |
|  Ehlers-Danlos Syndrome (EDS) (COL5A1, Labrador Retriever Variant) | Clear |



















OTHER RESULTS

| | |
|--|-------|
|  Enamel Hypoplasia (ENAM Deletion, Italian Greyhound Variant) | Clear |
|  Enamel Hypoplasia (ENAM SNP, Parson Russell Terrier Variant) | Clear |
|  Episodic Falling Syndrome (BCAN) | Clear |
|  Exercise-Induced Collapse, EIC (DNM1) | Clear |
|  Factor VII Deficiency (F7 Exon 5) | Clear |
|  Factor XI Deficiency (F11 Exon 7, Kerry Blue Terrier Variant) | Clear |
|  Familial Nephropathy (COL4A4 Exon 3, Cocker Spaniel Variant) | Clear |
|  Familial Nephropathy (COL4A4 Exon 30, English Springer Spaniel Variant) | Clear |
|  Fanconi Syndrome (FAN1, Basenji Variant) | Clear |
|  Fetal-Onset Neonatal Neuroaxonal Dystrophy (MFN2, Giant Schnauzer Variant) | Clear |
|  Glanzmann's Thrombasthenia Type I (ITGA2B Exon 13, Great Pyrenees Variant) | Clear |
|  Glanzmann's Thrombasthenia Type I (ITGA2B Exon 12, Otterhound Variant) | Clear |
|  Globoid Cell Leukodystrophy, Krabbe disease (GALC Exon 5, Terrier Variant) | Clear |
|  Glycogen Storage Disease Type IA, Von Gierke Disease, GSD IA (G6PC1, German Pinscher Variant) | Clear |
|  Glycogen Storage Disease Type IA, Von Gierke Disease, GSD IA (G6PC, Maltese Variant) | Clear |
|  Glycogen Storage Disease Type IIIA, GSD IIIA (AGL, Curly Coated Retriever Variant) | Clear |
|  Glycogen storage disease Type VII, Phosphofructokinase Deficiency, PFK Deficiency (PFKM, Whippet and English Springer Spaniel Variant) | Clear |
|  Glycogen storage disease Type VII, Phosphofructokinase Deficiency, PFK Deficiency (PFKM, Wachtelhund Variant) | Clear |

OTHER RESULTS

| | |
|--|-------|
| <input checked="" type="checkbox"/> GM1 Gangliosidosis (GLB1 Exon 2, Portuguese Water Dog Variant) | Clear |
| <input checked="" type="checkbox"/> GM1 Gangliosidosis (GLB1 Exon 15, Shiba Inu Variant) | Clear |
| <input checked="" type="checkbox"/> GM2 Gangliosidosis (HEXA, Japanese Chin Variant) | Clear |
| <input checked="" type="checkbox"/> GM2 Gangliosidosis (HEXB, Poodle Variant) | Clear |
| <input checked="" type="checkbox"/> Golden Retriever Progressive Retinal Atrophy 1, GR-PRA1 (SLC4A3) | Clear |
| <input checked="" type="checkbox"/> Golden Retriever Progressive Retinal Atrophy 2, GR-PRA2 (TTC8) | Clear |
| <input checked="" type="checkbox"/> Goniodysgenesis and Glaucoma, Pectinate Ligament Dysplasia, PLD (OLFM3) | Clear |
| <input checked="" type="checkbox"/> Hemophilia A (F8 Exon 11, German Shepherd Variant 1) | Clear |
| <input checked="" type="checkbox"/> Hemophilia A (F8 Exon 1, German Shepherd Variant 2) | Clear |
| <input checked="" type="checkbox"/> Hemophilia A (F8 Exon 10, Boxer Variant) | Clear |
| <input checked="" type="checkbox"/> Hemophilia B (F9 Exon 7, Terrier Variant) | Clear |
| <input checked="" type="checkbox"/> Hemophilia B (F9 Exon 7, Rhodesian Ridgeback Variant) | Clear |
| <input checked="" type="checkbox"/> Hereditary Ataxia (PNPLA8, Australian Shepherd Variant) | Clear |
| <input checked="" type="checkbox"/> Hereditary Ataxia, Cerebellar Degeneration (RAB24, Old English Sheepdog and Gordon Setter Variant) | Clear |
| <input checked="" type="checkbox"/> Hereditary Cataracts (HSF4 Exon 9, Australian Shepherd Variant) | Clear |
| <input checked="" type="checkbox"/> Hereditary Cataracts (FYCO1, Wirehaired Pointing Griffon Variant) | Clear |
| <input checked="" type="checkbox"/> Hereditary Cerebellar Ataxia (SELENOP, Belgian Shepherd Variant) | Clear |
| <input checked="" type="checkbox"/> Hereditary Footpad Hyperkeratosis (FAM83G, Terrier and Kromfohrlander Variant) | Clear |



















OTHER RESULTS

| | |
|--|-------|
|  Hereditary Footpad Hyperkeratosis (DSG1, Rottweiler Variant) | Clear |
|  Hereditary Nasal Parakeratosis (SUV39H2 Intron 4, Greyhound Variant) | Clear |
|  Hereditary Nasal Parakeratosis, HNPk (SUV39H2) | Clear |
|  Hypocatalasia, Acatlasemia (CAT) | Clear |
|  Hypomyelination and Tremors (FNIP2, Weimaraner Variant) | Clear |
|  Hypophosphatasia (ALPL Exon 9, Karelian Bear Dog Variant) | Clear |
|  Ichthyosis (NIPAL4, American Bulldog Variant) | Clear |
|  Ichthyosis (ASPRV1 Exon 2, German Shepherd Variant) | Clear |
|  Ichthyosis (SLC27A4, Great Dane Variant) | Clear |
|  Ichthyosis, Epidermolytic Hyperkeratosis (KRT10, Terrier Variant) | Clear |
|  Ichthyosis, ICH1 (PNPLA1, Golden Retriever Variant) | Clear |
|  Ichthyosis, ICH2 (ABHD5, Golden Retriever Variant) | Clear |
|  Inflammatory Myopathy (SLC25A12) | Clear |
|  Inherited Myopathy of Great Danes (BIN1) | Clear |
|  Inherited Selected Cobalamin Malabsorption with Proteinuria (CUBN, Komondor Variant) | Clear |
|  Intervertebral Disc Disease (Type I) (FGF4 retrogene - CFA12) | Clear |
|  Intestinal Lipid Malabsorption (ACSL5, Australian Kelpie) | Clear |
|  Junctional Epidermolysis Bullosa (LAMA3 Exon 66, Australian Cattle Dog Variant) | Clear |



















OTHER RESULTS

| | |
|---|-------|
| <input checked="" type="checkbox"/> Junctional Epidermolysis Bullosa (LAMB3 Exon 11, Australian Shepherd Variant) | Clear |
| <input checked="" type="checkbox"/> Juvenile Epilepsy (LGI2) | Clear |
| <input checked="" type="checkbox"/> Juvenile Laryngeal Paralysis and Polyneuropathy (RAB3GAP1, Rottweiler Variant) | Clear |
| <input checked="" type="checkbox"/> Juvenile Myoclonic Epilepsy (DIRAS1) | Clear |
| <input checked="" type="checkbox"/> L-2-Hydroxyglutaricaciduria, L2HGA (L2HGDH, Staffordshire Bull Terrier Variant) | Clear |
| <input checked="" type="checkbox"/> Lagotto Storage Disease (ATG4D) | Clear |
| <input checked="" type="checkbox"/> Laryngeal Paralysis (RAPGEF6, Miniature Bull Terrier Variant) | Clear |
| <input checked="" type="checkbox"/> Laryngeal Paralysis and Polyneuropathy (CNTNAP1, Leonberger, Saint Bernard, and Labrador Retriever variant) | Clear |
| <input checked="" type="checkbox"/> Late Onset Spinocerebellar Ataxia (CAPN1) | Clear |
| <input checked="" type="checkbox"/> Late-Onset Neuronal Ceroid Lipofuscinosis, NCL 12 (ATP13A2, Australian Cattle Dog Variant) | Clear |
| <input checked="" type="checkbox"/> Leonberger Polyneuropathy 1 (LPN1, ARHGEF10) | Clear |
| <input checked="" type="checkbox"/> Leonberger Polyneuropathy 2 (GJA9) | Clear |
| <input checked="" type="checkbox"/> Lethal Acrodermatitis, LAD (MKLN1) | Clear |
| <input checked="" type="checkbox"/> Leukodystrophy (TSEN54 Exon 5, Standard Schnauzer Variant) | Clear |
| <input checked="" type="checkbox"/> Ligneous Membranitis, LM (PLG) | Clear |
| <input checked="" type="checkbox"/> Limb Girdle Muscular Dystrophy (SGCD, Boston Terrier Variant) | Clear |
| <input checked="" type="checkbox"/> Limb-Girdle Muscular Dystrophy 2D (SGCA Exon 3, Miniature Dachshund Variant) | Clear |
| <input checked="" type="checkbox"/> Long QT Syndrome (KCNQ1) | Clear |



















OTHER RESULTS

| | |
|---|-------|
|  Lundehund Syndrome (LEPREL1) | Clear |
|  Macular Corneal Dystrophy, MCD (CHST6) | Clear |
|  Malignant Hyperthermia (RYR1) | Clear |
|  May-Hegglin Anomaly (MYH9) | Clear |
|  MDR1 Drug Sensitivity (ABCB1) | Clear |
|  Medium-Chain Acyl-CoA Dehydrogenase Deficiency, MCADD (ACADM, Cavalier King Charles Spaniel Variant) | Clear |
|  Methemoglobinemia (CYB5R3, Pit Bull Terrier Variant) | Clear |
|  Microphthalmia (RBP4 Exon 2, Soft Coated Wheaten Terrier Variant) | Clear |
|  Mucopolysaccharidosis IIIB, Sanfilippo Syndrome Type B, MPS IIIB (NAGLU, Schipperke Variant) | Clear |
|  Mucopolysaccharidosis Type IIIA, Sanfilippo Syndrome Type A, MPS IIIA (SGSH Exon 6, Dachshund Variant) | Clear |
|  Mucopolysaccharidosis Type IIIA, Sanfilippo Syndrome Type A, MPS IIIA (SGSH Exon 6, New Zealand Huntaway Variant) | Clear |
|  Mucopolysaccharidosis Type VI, Maroteaux-Lamy Syndrome, MPS VI (ARSB Exon 5, Miniature Pinscher Variant) | Clear |
|  Mucopolysaccharidosis Type VII, Sly Syndrome, MPS VII (GUSB Exon 3, German Shepherd Variant) | Clear |
|  Mucopolysaccharidosis Type VII, Sly Syndrome, MPS VII (GUSB Exon 5, Terrier Brasileiro Variant) | Clear |
|  Muscular Dystrophy (DMD, Cavalier King Charles Spaniel Variant 1) | Clear |
|  Muscular Dystrophy (DMD, Golden Retriever Variant) | Clear |
|  Muscular Dystrophy-Dystroglycanopathy (LARGE1, Labrador Retriever Variant) | Clear |
|  Musladin-Lueke Syndrome, MLS (ADAMTSL2) | Clear |



















OTHER RESULTS

| | |
|--|-------|
|  Myasthenia Gravis-Like Syndrome (CHRNE, Heiderterrier Variant) | Clear |
|  Myotonia Congenita (CLCN1 Exon 23, Australian Cattle Dog Variant) | Clear |
|  Myotonia Congenita (CLCN1 Exon 19, Labrador Retriever Variant) | Clear |
|  Myotonia Congenita (CLCN1 Exon 7, Miniature Schnauzer Variant) | Clear |
|  Narcolepsy (HCRTR2 Exon 1, Dachshund Variant) | Clear |
|  Narcolepsy (HCRTR2 Intron 4, Doberman Pinscher Variant) | Clear |
|  Narcolepsy (HCRTR2 Intron 6, Labrador Retriever Variant) | Clear |
|  Nemaline Myopathy (NEB, American Bulldog Variant) | Clear |
|  Neonatal Cerebellar Cortical Degeneration (SPTBN2, Beagle Variant) | Clear |
|  Neonatal Encephalopathy with Seizures, NEWS (ATF2) | Clear |
|  Neonatal Interstitial Lung Disease (LAMP3) | Clear |
|  Neuroaxonal Dystrophy, NAD (VPS11, Rottweiler Variant) | Clear |
|  Neuroaxonal Dystrophy, NAD (TECPR2, Spanish Water Dog Variant) | Clear |
|  Neuronal Ceroid Lipofuscinosis 1, NCL 1 (PPT1 Exon 8, Dachshund Variant 1) | Clear |
|  Neuronal Ceroid Lipofuscinosis 10, NCL 10 (CTSD Exon 5, American Bulldog Variant) | Clear |
|  Neuronal Ceroid Lipofuscinosis 2, NCL 2 (TPP1 Exon 4, Dachshund Variant 2) | Clear |
|  Neuronal Ceroid Lipofuscinosis 5, NCL 5 (CLN5 Exon 4 SNP, Border Collie Variant) | Clear |
|  Neuronal Ceroid Lipofuscinosis 5, NCL 5 (CLN5 Exon 4 Deletion, Golden Retriever Variant) | Clear |



















OTHER RESULTS

| | |
|--|-------|
|  Neuronal Ceroid Lipofuscinosis 6, NCL 6 (CLN6 Exon 7, Australian Shepherd Variant) | Clear |
|  Neuronal Ceroid Lipofuscinosis 7, NCL 7 (MFSD8, Chihuahua and Chinese Crested Variant) | Clear |
|  Neuronal Ceroid Lipofuscinosis 8, NCL 8 (CLN8, Australian Shepherd Variant) | Clear |
|  Neuronal Ceroid Lipofuscinosis 8, NCL 8 (CLN8 Exon 2, English Setter Variant) | Clear |
|  Neuronal Ceroid Lipofuscinosis 8, NCL 8 (CLN8 Insertion, Saluki Variant) | Clear |
|  Neuronal Ceroid Lipofuscinosis, Cerebellar Ataxia, NCL4A (ARSG Exon 2, American Staffordshire Terrier Variant) | Clear |
|  Oculocutaneous Albinism, OCA (SLC45A2 Exon 6, Bullmastiff Variant) | Clear |
|  Oculoskeletal Dysplasia 2 (COL9A2, Samoyed Variant) | Clear |
|  Osteochondrodysplasia (SLC13A1, Poodle Variant) | Clear |
|  Osteogenesis Imperfecta (COL1A2, Beagle Variant) | Clear |
|  Osteogenesis Imperfecta (SERPINH1, Dachshund Variant) | Clear |
|  Osteogenesis Imperfecta (COL1A1, Golden Retriever Variant) | Clear |
|  P2Y12 Receptor Platelet Disorder (P2Y12) | Clear |
|  Pachyonychia Congenita (KRT16, Dogue de Bordeaux Variant) | Clear |
|  Paroxysmal Dyskinesia, PxD (PIGN) | Clear |
|  Persistent Mullerian Duct Syndrome, PMDS (AMHR2) | Clear |
|  Pituitary Dwarfism (POU1F1 Intron 4, Karelian Bear Dog Variant) | Clear |
|  Platelet Factor X Receptor Deficiency, Scott Syndrome (TMEM16F) | Clear |

OTHER RESULTS

| | |
|---|-------|
|  Polycystic Kidney Disease, PKD (PKD1) | Clear |
|  Pompe's Disease (GAA, Finnish and Swedish Lapphund, Lapponian Herder Variant) | Clear |
|  Prekallikrein Deficiency (KLKB1 Exon 8) | Clear |
|  Primary Ciliary Dyskinesia, PCD (NME5, Alaskan Malamute Variant) | Clear |
|  Primary Ciliary Dyskinesia, PCD (STK36, Australian Shepherd Variant) | Clear |
|  Primary Ciliary Dyskinesia, PCD (CCDC39 Exon 3, Old English Sheepdog Variant) | Clear |
|  Primary Hyperoxaluria (AGXT) | Clear |
|  Primary Open Angle Glaucoma (ADAMTS17 Exon 11, Basset Fauve de Bretagne Variant) | Clear |
|  Primary Open Angle Glaucoma (ADAMTS10 Exon 17, Beagle Variant) | Clear |
|  Primary Open Angle Glaucoma (ADAMTS10 Exon 9, Norwegian Elkhound Variant) | Clear |
|  Primary Open Angle Glaucoma and Primary Lens Luxation (ADAMTS17 Exon 2, Chinese Shar-Pei Variant) | Clear |
|  Progressive Retinal Atrophy (SAG) | Clear |
|  Progressive Retinal Atrophy (IFT122 Exon 26, Lapponian Herder Variant) | Clear |
|  Progressive Retinal Atrophy, Bardet-Biedl Syndrome (BBS2 Exon 11, Shetland Sheepdog Variant) | Clear |
|  Progressive Retinal Atrophy, CNGA (CNGA1 Exon 9) | Clear |
|  Progressive Retinal Atrophy, crd1 (PDE6B, American Staffordshire Terrier Variant) | Clear |
|  Progressive Retinal Atrophy, crd4/cord1 (RPGRIP1) | Clear |
|  Progressive Retinal Atrophy, PRA1 (CNGB1) | Clear |

OTHER RESULTS

| | |
|--|-------|
|  Progressive Retinal Atrophy, PRA3 (FAM161A) | Clear |
|  Progressive Retinal Atrophy, rcd1 (PDE6B Exon 21, Irish Setter Variant) | Clear |
|  Proportionate Dwarfism (GH1 Exon 5, Chihuahua Variant) | Clear |
|  Protein Losing Nephropathy, PLN (NPHS1) | Clear |
|  Pyruvate Dehydrogenase Deficiency (PDP1, Spaniel Variant) | Clear |
|  Pyruvate Kinase Deficiency (PKLR Exon 5, Basenji Variant) | Clear |
|  Pyruvate Kinase Deficiency (PKLR Exon 7, Beagle Variant) | Clear |
|  Pyruvate Kinase Deficiency (PKLR Exon 10, Terrier Variant) | Clear |
|  Pyruvate Kinase Deficiency (PKLR Exon 7, Labrador Retriever Variant) | Clear |
|  Pyruvate Kinase Deficiency (PKLR Exon 7, Pug Variant) | Clear |
|  Raine Syndrome (FAM20C) | Clear |
|  Recurrent Inflammatory Pulmonary Disease, RIPD (AKNA, Rough Collie Variant) | Clear |
|  Renal Cystadenocarcinoma and Nodular Dermatofibrosis (FLCN Exon 7) | Clear |
|  Retina Dysplasia and/or Optic Nerve Hypoplasia (SIX6 Exon 1, Golden Retriever Variant) | Clear |
|  Sensory Neuropathy (FAM134B, Border Collie Variant) | Clear |
|  Severe Combined Immunodeficiency, SCID (PRKDC, Terrier Variant) | Clear |
|  Severe Combined Immunodeficiency, SCID (RAG1, Wetterhoun Variant) | Clear |
|  Shaking Puppy Syndrome (PLP1, English Springer Spaniel Variant) | Clear |

OTHER RESULTS

| | |
|---|-------|
| <input checked="" type="checkbox"/> Shar-Pei Autoinflammatory Disease, SPAID, Shar-Pei Fever (MTBP) | Clear |
| <input checked="" type="checkbox"/> Skeletal Dysplasia 2, SD2 (COL11A2, Labrador Retriever Variant) | Clear |
| <input checked="" type="checkbox"/> Skin Fragility Syndrome (PKP1, Chesapeake Bay Retriever Variant) | Clear |
| <input checked="" type="checkbox"/> Spinocerebellar Ataxia (SCN8A, Alpine Dachsbracke Variant) | Clear |
| <input checked="" type="checkbox"/> Spinocerebellar Ataxia with Myokymia and/or Seizures (KCNJ10) | Clear |
| <input checked="" type="checkbox"/> Spongy Degeneration with Cerebellar Ataxia 1 (KCNJ10) | Clear |
| <input checked="" type="checkbox"/> Spongy Degeneration with Cerebellar Ataxia 2 (ATP1B2) | Clear |
| <input checked="" type="checkbox"/> Stargardt Disease (ABCA4 Exon 28, Labrador Retriever Variant) | Clear |
| <input checked="" type="checkbox"/> Succinic Semialdehyde Dehydrogenase Deficiency (ALDH5A1 Exon 7, Saluki Variant) | Clear |
| <input checked="" type="checkbox"/> Thrombopathia (RASGRP1 Exon 5, Basset Hound Variant) | Clear |
| <input checked="" type="checkbox"/> Thrombopathia (RASGRP1 Exon 8, Landseer Variant) | Clear |
| <input checked="" type="checkbox"/> Trapped Neutrophil Syndrome, TNS (VPS13B) | Clear |
| <input checked="" type="checkbox"/> Ullrich-like Congenital Muscular Dystrophy (COL6A3 Exon 10, Labrador Retriever Variant) | Clear |
| <input checked="" type="checkbox"/> Ullrich-like Congenital Muscular Dystrophy (COL6A1 Exon 3, Landseer Variant) | Clear |
| <input checked="" type="checkbox"/> Unilateral Deafness and Vestibular Syndrome (PTPRQ Exon 39, Doberman Pinscher) | Clear |
| <input checked="" type="checkbox"/> Von Willebrand Disease Type I, Type I vWD (VWF) | Clear |
| <input checked="" type="checkbox"/> Von Willebrand Disease Type II, Type II vWD (VWF, Pointer Variant) | Clear |
| <input checked="" type="checkbox"/> Von Willebrand Disease Type III, Type III vWD (VWF Exon 4, Terrier Variant) | Clear |

OTHER RESULTS

| | |
|--|-----------|
| <input checked="" type="checkbox"/> Von Willebrand Disease Type III, Type III vWD (VWF Intron 16, Nederlandse Kooikerhondje Variant) | Clear |
| <input checked="" type="checkbox"/> Von Willebrand Disease Type III, Type III vWD (VWF Exon 7, Shetland Sheepdog Variant) | Clear |
| <input checked="" type="checkbox"/> X-Linked Hereditary Nephropathy, XLHN (COL4A5 Exon 35, Samoyed Variant 2) | Clear |
| <input checked="" type="checkbox"/> X-Linked Myotubular Myopathy (MTM1, Labrador Retriever Variant) | Clear |
| <input checked="" type="checkbox"/> X-linked Severe Combined Immunodeficiency, X-SCID (IL2RG Exon 1, Basset Hound Variant) | Clear |
| <input checked="" type="checkbox"/> X-linked Severe Combined Immunodeficiency, X-SCID (IL2RG, Corgi Variant) | Clear |
| <input checked="" type="checkbox"/> Xanthine Urolithiasis (XDH, Mixed Breed Variant) | Clear |
| <input checked="" type="checkbox"/> β -Mannosidosis (MANBA Exon 16, Mixed-Breed Variant) | Clear |
| Mast Cell Tumor | No result |

HEALTH REPORT

Notable result

ALT Activity

Zari inherited both copies of the variant we tested for Alanine Aminotransferase Activity

Why is this important to your vet?

Zari has two copies of a variant in the GPT gene and is likely to have a lower than average baseline ALT activity. ALT is a commonly used measure of liver health on routine veterinary blood chemistry panels. As such, your veterinarian may want to watch for changes in Zari's ALT activity above their current, healthy, ALT activity. As an increase above Zari's baseline ALT activity could be evidence of liver damage, even if it is within normal limits by standard ALT reference ranges.

What is Alanine Aminotransferase Activity?

ALT is a liver enzyme that vets measure to monitor liver health. With this result, your dog may naturally have a lower ALT baseline. Knowing this helps your veterinarian interpret future bloodwork results more accurately.

How vets diagnose this condition

Genetic testing is the only way to provide your veterinarian with this clinical tool.

How this condition is treated

Veterinarians may recommend blood work to establish a baseline ALT value for healthy dogs with one or two copies of this variant.

Actions to take if your dog is affected

- Talk to your vet about your dog's ALT result, as it may help them better interpret your dog's blood work.
- Dogs with this result do not exhibit symptoms or develop health issues associated with this variant.

HEALTH REPORT

Notable result

Degenerative Myelopathy, DM

Zari inherited one copy of the variant we tested for Degenerative Myelopathy, DM

What does this result mean?

This variant should not impact Zari's health. This variant is inherited in an autosomal recessive manner, meaning that a dog needs two copies of the variant to show signs of this condition. Zari is unlikely to develop this condition due to this variant because she only has one copy of the variant.

Impact on Breeding

This result is also important if you decide to breed this dog - to produce the healthiest puppies we recommend genetic testing any potential mates for this condition.

What is Degenerative Myelopathy, DM?

This condition affects the spinal cord nerves involved in movement, most noticeably in the hind limbs. It is progressive, meaning symptoms worsen over time, including weakness, muscle loss, and changes in walking.

When signs & symptoms develop in affected dogs

Affected dogs do not usually show signs of DM until they are at least 8 years old.

How vets diagnose this condition

Definitive diagnosis requires microscopic analysis of the spinal cord after death. However, veterinarians use clues such as genetic testing, breed, age, and other diagnostics to determine if DM is the most likely cause of your dog's clinical signs.

How this condition is treated

As dogs are seniors at the time of onset, the treatment for DM is aimed towards increasing their comfort through a combination of lifestyle changes, medication, and physical therapy.

Actions to take if your dog is affected

- Talk to your vet about your dog's degenerative myelopathy result, as it may influence how they monitor your dog's mobility and overall health, especially in their senior years.
- Keep your dog active with regular, low-impact exercise to help them maintain a healthy weight and support their mobility.
- Watch for changes in movement, such as wobbling, reluctance to jump, or dragging their back paws, and consult your vet if you notice any of these signs.
- Provide good traction in your home with rugs or mats to help prevent slipping as your dog ages. If mobility becomes difficult, ask your vet about supportive devices such as harnesses or wheelchairs.

INBREEDING AND DIVERSITY


CATEGORY

RESULT

Inbreeding | Gene: *n/a* | Genetic Result: **4%**

Inbreeding is a measure of how closely related this dog's parents were. The higher the number, the more closely related the parents. In general, greater inbreeding is associated with increased incidence of genetically inherited conditions.

4%




Immune Response 1 | Gene: *DRB1* | Genetic Result: **Low Diversity**

Diversity in the Major Histocompatibility Complex (MHC) region of the genome has been found in some studies to be associated with the incidence of certain autoimmune diseases. Dogs that have less diversity in the MHC region—i.e. the Dog Leukocyte Antigen (DLA) inherited from the mother is similar to the DLA inherited from the father—are considered less immunologically diverse. A High Diversity result means the dog has two highly dissimilar haplotypes. A Low Diversity result means the dog has two similar but not identical haplotypes. A No Diversity result means the dog has inherited identical haplotypes from both parents. Some studies have shown associations between certain DRB1 haplotypes and autoimmune diseases such as Cushing's disease, but these findings have yet to be scientifically validated.

Low Diversity

How common is this amount of diversity in mixed breed dogs:



Immune Response 2 | Gene: *DQA1 and DQB1* | Genetic Result: **High Diversity**

Diversity in the Major Histocompatibility Complex (MHC) region of the genome has been found in some studies to be associated with the incidence of certain autoimmune diseases. Dogs that have less diversity in the MHC region—i.e. the Dog Leukocyte Antigen (DLA) inherited from the mother is similar to the DLA inherited from the father—are considered less immunologically diverse. A High Diversity result means the dog has two highly dissimilar haplotypes. A Low Diversity result means the dog has two similar but not identical haplotypes. A No Diversity result means the dog has inherited identical haplotypes from both parents. A number of studies have shown correlations of DQA-DQB1 haplotypes and certain autoimmune diseases; however, these have not yet been scientifically validated.

High Diversity

How common is this amount of diversity in mixed breed dogs:

